

Table 4F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 1
Run Date: 9 / 21 / 05

Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Local Agency							
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
3 Rivers	1,626	89.1	1.6	9.2	0.1	0.1	0.0
Allen	690	98.4	0.7	0.4	0.0	0.4	0.0
Anderson	427	91.6	4.4	4.0	0.0	0.0	0.0
Barren Riv	7,291	81.2	9.7	6.1	0.4	2.6	0.0
Bourbon	530	74.3	9.8	14.9	0.4	0.6	0.0
Boyd	1,593	97.0	1.9	0.9	0.1	0.1	0.0
Boyle	702	93.6	6.4	0.0	0.0	0.0	0.0
Bracken	290	98.3	0.3	1.4	0.0	0.0	0.0
Breathitt	698	99.6	0.3	0.1	0.0	0.0	0.0
Breckinrid	666	94.9	4.7	0.3	0.0	0.2	0.0
Buffalo Tr	701	90.9	5.7	2.0	0.3	1.1	0.0
Bullitt	1,352	97.9	1.2	0.8	0.0	0.1	0.0
C & Y	1,392	4.0	92.3	0.0	1.1	2.6	0.0
Christian	5,147	66.5	27.2	4.6	0.2	1.4	0.0
Clark	1,004	88.7	6.4	3.5	0.3	1.1	0.0
Cumb Val	6,617	98.4	1.4	0.1	0.1	0.1	0.0
Estill	692	99.6	0.3	0.0	0.0	0.1	0.0
Fayette	5,967	43.5	30.2	21.8	0.6	3.9	0.0
Fleming	642	97.2	1.6	1.2	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 9 / 21 / 05

Comparison of Racial and Ethnic Distribution by Local Agency							
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Floyd	2,118	99.4	0.5	0.1	0.0	0.0	0.0
Franklin	1,228	76.8	14.2	7.9	0.2	0.9	0.0
Garrard	590	95.1	1.2	3.7	0.0	0.0	0.0
Gateway	2,173	98.8	0.5	0.5	0.1	0.1	0.0
Green Riv	7,283	86.7	8.7	3.8	0.1	0.8	0.0
Greenup	1,094	99.3	0.6	0.1	0.0	0.0	0.0
Hopkins	1,666	85.7	11.4	2.1	0.2	0.6	0.0
Jefferson	13,470	54.7	35.0	7.0	0.3	2.9	0.0
Jessamine	1,293	90.3	5.3	3.2	0.2	1.1	0.0
Johnson	1,189	99.9	0.1	0.0	0.0	0.0	0.0
Knox	1,640	99.2	0.7	0.1	0.0	0.0	0.0
KY River	5,107	99.2	0.7	0.1	0.0	0.1	0.0
Lake Cumb	8,511	94.4	1.9	3.5	0.0	0.2	0.0
Laurel	1,989	98.8	0.5	0.2	0.3	0.3	0.0
Lawrence	733	99.6	0.4	0.0	0.0	0.0	0.0
Lewis	617	99.8	0.0	0.0	0.0	0.2	0.0
Lincoln	919	94.0	2.9	2.9	0.1	0.0	0.0
Lincoln Tr	8,199	84.9	11.1	3.2	0.2	0.6	0.0
Little San	1,810	99.2	0.3	0.2	0.1	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 3
Run Date: 9 / 21 / 05

Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Local Agency							
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Madison	1,985	89.1	7.5	2.1	0.3	1.1	0.0
Magoffin	862	99.7	0.0	0.3	0.0	0.0	0.0
Marshall	962	99.3	0.7	0.0	0.0	0.0	0.0
Martin	775	99.7	0.3	0.0	0.0	0.0	0.0
Mercer	716	90.5	3.2	5.9	0.1	0.3	0.0
Monroe	452	92.0	4.4	3.5	0.0	0.0	0.0
Montgomery	1,016	96.1	1.0	2.4	0.0	0.6	0.0
Muhlenberg	1,099	94.4	5.1	0.5	0.0	0.1	0.0
N Central	2,026	69.4	5.6	25.0	0.0	0.0	0.0
N KY	7,176	88.6	6.8	4.0	0.1	0.4	0.0
Oldham	832	80.2	4.2	15.3	0.0	0.4	0.0
Park DuV	1,486	11.3	74.4	11.1	0.2	3.0	0.0
Pennyrile	2,624	91.2	6.0	1.7	0.2	0.9	0.0
Pike	2,218	99.1	0.5	0.1	0.1	0.2	0.0
Powell	774	97.0	1.0	1.8	0.0	0.1	0.0
Purchase	4,588	68.0	23.8	7.0	0.2	1.1	0.0
Todd	569	72.6	14.1	2.8	0.4	10.2	0.0
WEDCO	1,948	88.1	3.9	7.4	0.3	0.4	0.0
Whitley	2,685	99.7	0.1	0.0	0.0	0.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4F
2004 Pediatric Nutrition Surveillance ⁽¹⁾

Page: 4
Run Date: 9 / 21 / 05

Kentucky
Comparison of Racial and Ethnic Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Local Agency							
Local Agency	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Woodford	650	68.3	8.6	23.1	0.0	0.0	0.0
Unknown Local Agencies	998	92.4	5.1	2.0	0.0	0.5	0.0
Kentucky	136,097	82.3	11.9	4.6	0.2	1.0	0.0
Nation (Prior Year)	5,091,536	38.8	20.4	32.2	1.3	3.6	3.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 9 / 21 / 05

Comparison of Age Distribution by Local Agency				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
3 Rivers	1,626	35.1	23.9	41.0
Allen	690	28.7	23.2	48.1
Anderson	427	38.9	21.3	39.8
Barren Riv	7,291	38.5	23.5	38.0
Bourbon	530	32.1	26.6	41.3
Boyd	1,593	35.7	27.0	37.3
Boyle	702	35.0	26.4	38.6
Bracken	290	35.5	26.9	37.6
Breathitt	698	23.8	28.1	48.1
Breckinrid	666	34.4	24.8	40.8
Buffalo Tr	701	33.7	23.5	42.8
Bullitt	1,352	36.8	27.8	35.4
C & Y	1,392	33.1	26.5	40.4
Christian	5,147	33.9	22.0	44.1
Clark	1,004	39.8	23.6	36.6
Cumb Val	6,617	28.8	24.6	46.6
Estill	692	29.9	26.6	43.5
Fayette	5,967	37.9	23.6	38.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 9 / 21 / 05

Comparison of Age Distribution by Local Agency				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Fleming	642	28.5	24.0	47.5
Floyd	2,118	29.8	24.4	45.8
Franklin	1,228	46.2	19.5	34.4
Garrard	590	27.1	27.5	45.4
Gateway	2,173	28.7	25.6	45.7
Green Riv	7,283	36.7	22.8	40.6
Greenup	1,094	30.2	24.2	45.6
Hopkins	1,666	40.5	19.4	40.1
Jefferson	13,470	38.2	24.2	37.6
Jessamine	1,293	38.1	22.0	40.0
Johnson	1,189	29.1	24.7	46.2
Knox	1,640	29.4	25.9	44.7
KY River	5,107	33.1	22.6	44.3
Lake Cumb	8,511	32.1	24.4	43.5
Laurel	1,989	37.6	23.1	39.3
Lawrence	733	33.3	23.1	43.7
Lewis	617	34.4	21.1	44.6
Lincoln	919	37.1	19.2	43.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 9 / 21 / 05

Comparison of Age Distribution by Local Agency				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Lincoln Tr	8,199	32.8	24.0	43.2
Little San	1,810	27.2	25.0	47.8
Madison	1,985	36.3	24.5	39.2
Magoffin	862	28.0	23.0	49.1
Marshall	962	33.2	23.9	42.9
Martin	775	29.2	23.9	47.0
Mercer	716	34.9	22.8	42.3
Monroe	452	37.4	20.4	42.3
Montgomery	1,016	33.1	22.0	44.9
Muhlenberg	1,099	32.6	25.1	42.3
N Central	2,026	32.9	27.0	40.1
N KY	7,176	31.9	26.7	41.3
Oldham	832	30.8	25.6	43.6
Park DuV	1,486	36.7	23.3	40.0
Pennyrile	2,624	29.9	28.0	42.1
Pike	2,218	36.7	22.8	40.5
Powell	774	27.3	23.6	49.1
Purchase	4,588	31.3	26.8	42.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Age Distribution by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 9 / 21 / 05

Comparison of Age Distribution by Local Agency				
Local Agency	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Todd	569	33.4	23.2	43.4
WEDCO	1,948	36.8	24.6	38.7
Whitley	2,685	34.6	22.5	42.9
Woodford	650	32.0	27.5	40.5
Unknown Local Agen	998	100.0	0.0	0.0
Kentucky	136,097	34.6	24.0	41.4
Nation (Prior Year)	5,091,536	36.5	21.5	41.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Local Agency (3,4)											
Local Agency	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
3 Rivers	516	7.0 (4)	7.2 (39)	1,575	7.7(28)	5.1(38)	13.6 (6)	645	19.4(41)	15.8(16)	1,056 8.6 (19)
Allen	176	7.4 (7)	5.7 (15)	673	6.7(20)	3.6(19)	16.5(25)	323	22.9(56)	16.4(20)	379 8.2 (16)
Anderson	145	15.2 (57)	7.6 (44)	407	7.9(30)	3.9(23)	18.7(43)	163	22.1(55)	22.1(51)	272 11.4 (29)
Barren Riv	2,341	8.1 (14)	8.7 (51)	7,016	5.8(13)	3.9(23)	16.8(28)	2,696	19.6(43)	20.1(39)	4,271 13.6 (40)
Bourbon	159	10.7 (38)	8.2 (48)	504	8.3(36)	3.2(12)	19.8(47)	214	18.2(30)	22.0(49)	362 27.1 (58)
Boyd	513	12.3 (50)	6.0 (17)	1,541	6.2(15)	7.3(58)	15.2(18)	582	15.3(11)	16.2(19)	967 15.0 (43)
Boyle	225	11.1 (41)	5.3 (12)	668	4.9 (7)	5.8(48)	18.3(37)	260	20.0(46)	16.5(21)	419 8.1 (15)
Bracken	96	* *	* *	285	4.9 (7)	5.3(42)	21.8(52)	109	17.4(21)	31.2(58)	186 6.5 (7)
Breathitt	156	4.5 (1)	7.7 (45)	663	8.3(36)	2.6 (7)	22.2(55)	329	18.5(34)	20.1(39)	493 7.9 (13)
Breckinrid	194	8.8 (20)	6.2 (18)	650	8.8(42)	2.3 (2)	14.9(14)	273	18.7(35)	14.7(11)	456 5.0 (2)
Buffalo Tr	221	7.7 (8)	13.1 (58)	678	7.5(26)	5.2(40)	18.3(37)	297	16.2(14)	17.5(28)	428 7.9 (13)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Page: 2
Run Date: 9 / 21 / 05

Comparison of Growth and Anemia Indicators by Local Agency (3,4)											
Local Agency	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Bullitt	454	11.7 (44)	7.0 (33)	1,295	8.5(40)	2.5 (5)	15.1(17)	461	16.1(13)	13.2 (5)	768 15.6 (46)
C & Y	396	13.6 (55)	4.3 (6)	1,337	8.5(40)	5.5(44)	17.9(34)	557	20.1(47)	20.1(39)	210 19.0 (54)
Christian	1,149	8.0 (11)	6.3 (21)	4,967	4.8 (5)	6.5(57)	11.1 (2)	2,222	12.0 (2)	7.8 (1)	3,534 22.7 (57)
Clark	359	10.6 (35)	6.7 (26)	965	4.1 (3)	3.4(14)	12.0 (3)	356	16.6(19)	17.1(27)	597 13.1 (39)
Cumb Val	1,762	10.6 (35)	6.5 (24)	6,329	8.3(36)	5.9(51)	19.9(48)	3,003	18.9(36)	20.6(43)	4,330 5.9 (4)
Estill	183	11.5 (43)	4.4 (7)	673	10.4(55)	3.4(14)	18.6(41)	288	20.1(47)	19.8(36)	488 11.9 (31)
Fayette	1,903	10.7 (38)	6.7 (26)	5,691	8.2(35)	4.0(25)	16.9(29)	2,248	21.0(50)	20.0(37)	3,414 16.5 (50)
Fleming	170	5.9 (2)	11.2 (55)	615	6.2(15)	3.4(14)	18.2(36)	297	19.9(45)	14.1 (8)	441 16.3 (48)
Floyd	588	13.9 (56)	4.8 (9)	2,007	5.6(12)	6.1(55)	12.7 (5)	928	14.7 (9)	12.5 (4)	1,418 7.8 (12)
Franklin	463	8.9 (22)	3.9 (2)	1,180	6.7(20)	5.1(38)	18.6(41)	415	16.4(16)	21.9(48)	723 18.1 (52)
Garrard	136	11.8 (46)	5.1 (11)	575	6.8(22)	3.8(22)	15.5(21)	267	15.4(12)	15.7(14)	370 17.0 (51)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Page: 3
Run Date: 9 / 21 / 05

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)														
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)			
	Low		High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)		Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)		Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Gateway	551	9.3 (27)	12.2 (57)		2,095	6.4 (18)	4.2 (31)	15.8 (22)		969	17.4 (21)	16.8 (24)	1,447	9.2 (24)
Green Riv	2,201	9.2 (25)	5.6 (14)		7,027	10.2 (54)	3.0 (11)	18.5 (39)		2,865	19.2 (40)	20.5 (42)	4,374	8.6 (19)
Greenup	281	8.2 (16)	8.9 (52)		1,067	5.2 (10)	5.8 (48)	17.9 (34)		491	16.3 (15)	20.0 (37)	693	6.3 (6)
Hopkins	530	7.7 (8)	6.6 (25)		1,596	12.3 (58)	2.3 (2)	19.7 (46)		654	21.3 (52)	20.8 (45)	1,057	9.9 (27)
Jefferson	4,303	9.6 (30)	6.4 (23)		12,750	7.1 (25)	4.8 (35)	15.3 (20)		4,900	16.9 (20)	14.1 (8)	8,272	12.9 (36)
Jessamine	398	10.3 (33)	8.0 (46)		1,243	7.9 (30)	4.0 (25)	12.6 (4)		508	14.6 (8)	13.2 (5)	792	13.0 (38)
Johnson	323	7.1 (6)	5.9 (16)		1,141	6.6 (19)	4.6 (34)	16.4 (24)		535	17.8 (27)	16.8 (24)	786	19.1 (55)
Knox	429	10.3 (33)	7.0 (33)		1,570	4.6 (4)	5.8 (48)	17.5 (32)		710	17.5 (23)	23.8 (55)	1,053	16.0 (47)
KY River	1,428	9.2 (25)	6.7 (26)		4,827	9.9 (51)	2.5 (5)	26.7 (58)		2,201	19.8 (44)	21.1 (46)	3,471	8.8 (21)
Lake Cumb	2,436	8.0 (11)	7.1 (36)		8,214	6.3 (17)	4.5 (33)	15.0 (16)		3,600	18.1 (29)	16.5 (21)	5,170	8.9 (23)
Laurel	619	9.0 (24)	7.3 (41)		1,901	6.8 (22)	5.3 (42)	20.0 (50)		759	22.9 (56)	23.8 (55)	1,183	12.7 (35)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Page: 4
Run Date: 9 / 21 / 05

Comparison of Growth and Anemia Indicators by Local Agency (3,4)											
Local Agency	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Lawrence	220	19.5 (58)	4.1 (4)	680	9.9(51)	5.6(47)	19.4(45)	309	20.4(49)	22.0(49)	451 7.5 (10)
Lewis	193	9.3 (27)	11.9 (56)	582	6.9(24)	3.6(19)	14.8(12)	268	14.2 (7)	15.7(14)	351 2.8 (1)
Lincoln	273	11.4 (42)	5.5 (13)	883	7.8(29)	5.9(51)	22.2(55)	391	21.0(50)	22.3(52)	630 8.4 (18)
Lincoln Tr	2,198	8.7 (19)	6.2 (18)	7,913	9.6(47)	3.5(18)	16.6(26)	3,430	19.1(38)	14.7(11)	5,674 8.8 (21)
Little San	463	9.9 (31)	6.7 (26)	1,762	4.8 (5)	5.5(44)	14.6(10)	847	15.0(10)	16.8(24)	1,160 9.2 (24)
Madison	620	12.1 (48)	8.1 (47)	1,892	5.9(14)	6.0(53)	13.6 (6)	764	18.3(33)	19.2(34)	1,283 11.4 (29)
Magoffin	223	8.1 (14)	9.0 (53)	818	8.1(32)	3.4(14)	24.3(57)	406	21.7(53)	22.7(54)	542 16.4 (49)
Marshall	286	7.0 (4)	9.1 (54)	922	9.8(49)	6.0(53)	14.4 (9)	394	12.7 (4)	10.2 (2)	618 15.0 (43)
Martin	218	10.6 (35)	7.3 (41)	751	8.4(39)	2.7 (8)	22.0(54)	352	18.2(30)	20.7(44)	549 7.7 (11)
Mercer	204	7.8 (10)	4.4 (7)	694	10.1(53)	3.3(13)	14.0 (8)	289	13.5 (6)	16.6(23)	473 6.1 (5)
Monroe	145	6.9 (3)	6.2 (18)	441	9.8(49)	3.6(19)	16.1(23)	183	17.5(23)	17.5(28)	278 12.6 (34)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)														
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)			
	Low		High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)		Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)		Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Montgomery	282	11.0 (40)	8.5 (49)		969	9.6(47)	2.3 (2)	14.7(11)		440	17.5(23)	13.6 (7)	714	8.3 (17)
Muhlenberg	338	8.9 (22)	6.8 (31)		1,055	5.0 (9)	4.1(28)	15.2(18)		463	16.4(16)	19.2(34)	628	12.9 (36)
N Central	589	8.0 (11)	7.0 (33)		1,912	14.6(59)	2.8 (9)	29.4(59)		775	21.9(54)	24.3(57)	1,318	14.6 (42)
N KY	1,895	11.7 (44)	7.4 (43)		6,788	9.2(45)	4.1(28)	18.5(39)		2,890	18.0(28)	19.1(33)	4,633	12.4 (32)
Oldham	220	11.8 (46)	6.8 (31)		801	12.0(57)	2.9(10)	21.8(52)		350	17.7(26)	18.9(32)	496	12.5 (33)
Park DuV	461	13.4 (53)	3.9 (2)		1,407	9.2(45)	4.1(28)	14.9(14)		572	18.2(30)	16.1(18)	1,001	9.4 (26)
Pennyrile	718	12.5 (51)	6.7 (26)		2,518	7.5(26)	4.0(25)	17.3(30)		1,077	18.9(36)	15.9(17)	1,739	15.5 (45)
Pike	672	10.0 (32)	7.1 (36)		2,100	4.0 (2)	6.4(56)	16.6(26)		868	16.5(18)	18.4(31)	1,352	7.2 (9)
Powell	199	13.1 (52)	3.5 (1)		734	8.9(44)	2.2 (1)	19.9(48)		370	23.2(58)	21.6(47)	565	5.7 (3)
Purchase	1,358	8.3 (17)	7.1 (36)		4,342	8.1(32)	4.2(31)	20.2(51)		1,862	19.5(42)	17.6(30)	2,911	19.6 (56)
Todd	171	13.5 (54)	4.1 (4)		539	11.7(56)	5.2(40)	17.8(33)		240	13.3 (5)	12.1 (3)	375	18.7 (53)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Local Agency (3,4)															
Local Agency	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)				
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)								
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)		
WEDCO	633	12.2 (49)	4.9 (10)	1,837	8.1 (32)	4.8 (35)	17.3 (30)	737	12.5 (3)	15.3 (13)		1,205	11.1 (28)		
Whitley	763	9.3 (27)	7.2 (39)	2,591	8.8 (42)	4.8 (35)	18.8 (44)	1,119	19.1 (38)	22.3 (52)		1,804	6.9 (8)		
Woodford	165	8.5 (18)	8.5 (49)	636	5.3 (11)	5.5 (44)	14.8 (12)	261	10.7 (1)	14.2 (10)		379	14.5 (41)		
Unknown Local Agenc	997	8.8 (20)	6.3 (21)	873	0.7 (1)	9.7 (59)	3.2 (1)	0	*	*	*	*	0	*	*
Kentucky	40,308	9.7	6.8	130,165	7.7	4.4	17.2	54,782	18.1	17.6		85,009	11.8		
Nation (Prior Year)	1,172,866	9.1	7.3	4,838,257	6.2	5.2	13.6	2,044,705	15.7	14.7		3,118,041	12.8		

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 1
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
3 Rivers	518	20.3 (16)	133	6.0 (13)	276	2.9 (33)
Allen	176	19.9 (17)	52	* *	128	2.3 (41)
Anderson	149	24.8 (6)	40	* *	72	* *
Barren Riv	2,393	19.7 (20)	669	6.6 (11)	1,165	2.9 (33)
Bourbon	159	20.8 (13)	41	* *	107	4.7 (10)
Boyd	513	12.1 (43)	161	1.9 (30)	295	2.4 (39)
Boyle	228	10.5 (51)	79	* *	135	3.7 (20)
Bracken	96	* *	26	* *	55	* *
Breathitt	156	9.6 (53)	75	* *	137	2.9 (33)
Breckinrid	196	19.9 (17)	59	* *	106	1.9 (46)
Buffalo Tr	222	16.7 (28)	48	* *	105	0.0 (54)
Bullitt	456	17.5 (24)	134	3.0 (25)	277	2.9 (33)
C & Y	397	9.6 (53)	148	6.8 (10)	248	2.0 (44)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 2
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Christian	1,194	24.2 (9)	545	9.2 (5)	680	4.6 (11)
Clark	360	20.8 (13)	79	* *	156	1.9 (46)
Cumb Val	1,781	11.1 (47)	573	3.0 (25)	1,127	2.6 (37)
Estill	184	10.3 (52)	69	* *	123	1.6 (49)
Fayette	1,909	38.1 (1)	605	13.6 (1)	978	5.2 (6)
Fleming	173	16.8 (26)	48	* *	111	4.5 (12)
Floyd	588	8.2 (57)	197	3.6 (22)	378	3.4 (25)
Franklin	463	27.0 (4)	115	9.6 (4)	155	5.2 (6)
Garrard	138	10.9 (49)	60	* *	111	3.6 (22)
Gateway	554	11.6 (44)	211	4.3 (19)	402	3.2 (30)
Green Riv	2,202	16.3 (31)	704	3.6 (22)	1,148	2.0 (44)
Greenup	283	14.5 (36)	103	1.9 (30)	173	4.0 (17)
Hopkins	547	16.6 (30)	149	4.0 (21)	212	3.3 (29)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 3
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Jefferson	4,352	24.8 (6)	1,361	8.4 (6)	2,263	4.9 (9)
Jessamine	400	22.3 (11)	146	10.3 (2)	207	4.3 (14)
Johnson	323	15.8 (33)	98	* *	201	3.5 (23)
Knox	442	9.0 (55)	161	0.0 (32)	303	1.0 (52)
KY River	1,467	12.3 (42)	396	3.0 (25)	755	2.3 (41)
Lake Cumb	2,450	13.4 (40)	783	4.1 (20)	1,413	2.6 (37)
Laurel	630	14.3 (38)	182	2.2 (29)	313	2.2 (43)
Lawrence	220	8.2 (57)	67	* *	118	3.4 (25)
Lewis	193	16.1 (32)	43	* *	102	0.0 (54)
Lincoln	273	15.4 (34)	80	* *	114	1.8 (48)
Lincoln Tr	2,206	16.9 (25)	788	6.0 (13)	1,366	5.5 (5)
Little San	463	18.4 (23)	150	4.7 (17)	296	3.4 (25)
Madison	625	19.2 (21)	186	10.2 (3)	336	5.1 (8)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 4
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Magoffin	223	10.8 (50)	86	* *	151	1.3 (51)
Marshall	289	11.4 (45)	92	* *	156	3.2 (30)
Martin	219	8.7 (56)	55	* *	123	0.8 (53)
Mercer	204	11.3 (46)	70	* *	110	10.0 (1)
Monroe	145	14.5 (36)	34	* *	64	* *
Montgomery	282	16.7 (28)	104	7.7 (7)	151	4.0 (17)
Muhlenberg	338	13.3 (41)	94	* *	194	1.5 (50)
N Central	590	19.8 (19)	199	4.5 (18)	387	3.4 (25)
N KY	1,922	16.8 (26)	734	6.5 (12)	1,300	4.1 (15)
Oldham	224	30.8 (2)	89	* *	151	9.9 (2)
Park DuV	461	20.4 (15)	125	7.2 (8)	233	6.4 (4)
Pennyrile	719	25.7 (5)	265	6.0 (13)	533	4.1 (15)
Pike	677	11.1 (47)	212	3.3 (24)	322	3.1 (32)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2004 Pediatric Nutrition Surveillance (1)
Kentucky

Page: 5
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Powell	199	15.1 (35)	64	* *	127	2.4 (39)
Purchase	1,360	18.8 (22)	405	7.2 (8)	827	4.5 (12)
Todd	171	21.6 (12)	40	* *	103	6.8 (3)
WEDCO	634	22.4 (10)	167	5.4 (16)	349	4.0 (17)
Whitley	763	13.6 (39)	229	2.6 (28)	408	3.7 (20)
Woodford	165	27.9 (3)	77	* *	114	3.5 (23)
Unknown Local Agencies	998	24.8 (6)	0	* *	0	* *
Kentucky	40,662	18.5	12,705	5.8	22,450	3.6
Nation (Prior Year)	928,626	53.2	490,960	21.5	504,421	13.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 9 / 21 / 05

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)														
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)			
	Low		High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)		Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)		Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
3 Rivers	1,156	8.3 (7)	6.7 (20)		4,265	8.1 (40)	4.5(34)	15.6(20)		1,896	19.4(44)	17.0(26)	3,068	9.2 (22)
Allen	436	8.7 (15)	8.7 (46)		1,946	6.7(19)	3.0 (9)	16.9(29)		1,011	21.9(58)	17.5(28)	1,119	9.0 (20)
Anderson	300	11.3 (50)	6.7 (20)		1,081	8.1 (40)	4.2(31)	20.4(51)		492	20.5(54)	22.0(54)	802	9.2 (22)
Barren Riv	5,623	8.8 (17)	7.8 (40)		19,566	6.3(14)	3.7(21)	16.8(28)		8,244	18.8(35)	18.6(36)	11,847	12.6 (38)
Bourbon	384	10.9 (45)	8.3 (44)		1,470	8.9(48)	2.5 (5)	20.5(53)		655	18.6(31)	22.3(55)	1,098	27.6 (59)
Boyd	1,281	12.3 (55)	7.0 (27)		4,216	6.5(18)	7.9(58)	14.7 (9)		1,769	17.6(22)	16.1(22)	2,604	12.2 (36)
Boyle	553	11.0 (46)	6.7 (20)		1,790	6.1(11)	5.1(42)	17.8(37)		724	18.0(27)	17.7(31)	1,105	8.1 (14)
Bracken	217	7.4 (3)	6.5 (15)		761	8.5(46)	5.9(50)	23.8(56)		343	15.7(10)	30.9(58)	513	6.0 (3)
Breathitt	436	7.8 (4)	8.5 (45)		1,970	6.9(22)	3.5(19)	21.0(54)		1,064	17.5(20)	19.5(41)	1,425	7.7 (12)
Breckinrid	420	8.6 (14)	5.7 (8)		1,756	7.6(30)	2.7 (7)	15.4(17)		833	19.4(44)	14.2 (8)	1,315	3.7 (1)
Buffalo Tr	515	8.0 (5)	10.3 (57)		2,000	8.4(43)	5.3(45)	18.1(39)		958	17.3(18)	15.8(18)	1,359	9.9 (25)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 9 / 21 / 05

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)													
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Bullitt	1,129	9.0 (23)	8.8 (48)	3,503	8.0(38)	3.4(16)	16.4(25)	1,374	15.7(10)	15.4(14)		1,965	18.3 (54)
C & Y	1,117	12.4 (57)	5.5 (5)	3,757	8.0(38)	5.2(43)	17.5(33)	1,646	19.1(40)	20.1(46)		767	18.9 (55)
Christian	3,296	8.5 (11)	7.6 (37)	14,371	5.3 (8)	6.4(54)	10.9 (2)	6,567	12.5 (3)	8.6 (1)		9,485	25.3 (58)
Clark	806	11.9 (54)	6.5 (15)	2,540	5.5(10)	3.7(21)	13.8 (4)	1,028	17.7(23)	17.0(26)		1,552	12.9 (39)
Cumb Val	4,038	9.9 (31)	7.4 (33)	17,930	7.0(25)	5.3(45)	19.6(47)	9,265	17.7(23)	19.1(38)		12,510	7.3 (10)
Estill	455	8.1 (6)	6.6 (17)	1,877	6.9(22)	3.2(12)	17.6(34)	900	19.1(40)	19.9(43)		1,429	10.4 (28)
Fayette	4,776	11.2 (49)	6.9 (26)	15,631	7.6(30)	3.9(24)	16.1(24)	6,365	20.0(50)	18.1(34)		9,299	15.4 (48)
Fleming	386	6.2 (1)	10.4 (58)	1,765	6.2(13)	4.0(26)	16.9(29)	927	15.9(12)	15.5(15)		1,377	16.4 (50)
Floyd	1,380	11.3 (50)	6.4 (14)	5,724	5.0 (6)	7.5(57)	14.0 (5)	2,868	12.9 (4)	13.8 (7)		4,057	8.4 (17)
Franklin	1,028	8.9 (19)	5.5 (5)	3,086	7.8(35)	6.0(51)	18.4(43)	1,157	18.2(30)	21.3(52)		2,100	17.1 (51)
Garrard	360	9.2 (27)	5.8 (10)	1,676	6.9(22)	3.3(14)	14.7 (9)	820	18.9(37)	14.4 (9)		992	15.4 (48)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 9 / 21 / 05

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)														
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)			
	Low		High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)		Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)		Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Gateway	1,455	8.5 (11)	11.7 (59)		6,532	6.3(14)	3.7(21)	17.8(37)		3,277	17.8(26)	18.0(33)	4,401	11.6 (33)
Green Riv	5,208	9.5 (29)	5.8 (10)		19,400	10.1(55)	2.9 (8)	18.6(44)		8,692	18.6(31)	19.2(40)	12,298	9.3 (24)
Greenup	748	10.0 (35)	7.9 (41)		3,028	4.0 (3)	7.2(56)	15.2(12)		1,435	17.5(20)	16.0(21)	1,939	7.0 (8)
Hopkins	1,165	10.3 (40)	5.8 (10)		4,449	13.1(58)	1.7 (1)	20.1(49)		2,031	21.2(55)	19.1(38)	3,241	10.9 (31)
Jefferson	11,412	9.9 (31)	6.6 (17)		35,969	7.7(32)	4.7(37)	15.7(21)		14,474	17.0(17)	15.1(13)	23,920	11.7 (35)
Jessamine	885	8.9 (19)	9.0 (50)		3,327	8.9(48)	3.4(16)	15.7(21)		1,470	16.9(16)	14.8(11)	2,229	10.6 (29)
Johnson	739	8.5 (11)	7.2 (29)		3,282	5.3 (8)	5.3(45)	17.0(31)		1,703	16.4(15)	16.6(24)	2,360	20.0 (56)
Knox	1,033	11.4 (52)	7.6 (37)		4,585	5.0 (6)	4.1(30)	18.1(39)		2,341	19.1(40)	21.7(53)	3,203	13.3 (42)
KY River	3,319	9.1 (24)	6.6 (17)		13,775	9.4(52)	2.4 (4)	26.4(58)		6,903	18.6(31)	21.1(50)	10,603	8.8 (18)
Lake Cumb	5,693	8.9 (19)	7.4 (33)		22,854	6.8(20)	4.0(26)	15.3(16)		10,755	18.0(27)	16.5(23)	13,790	8.9 (19)
Laurel	1,479	9.9 (31)	7.6 (37)		5,053	7.0(25)	5.2(43)	17.7(36)		2,124	21.6(57)	21.0(49)	2,954	14.0 (44)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)														
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)			
	Low		High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)		Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)		Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lawrence	481	16.2 (59)	5.0 (4)		1,883	7.3(27)	5.0(40)	19.4(46)		929	21.2(55)	19.9(43)	1,246	6.9 (7)
Lewis	403	8.9 (19)	9.9 (56)		1,643	6.1(11)	3.5(19)	15.2(12)		830	19.2(43)	15.9(19)	950	4.1 (2)
Lincoln	655	11.5 (53)	7.3 (30)		2,590	8.5(46)	4.8(39)	21.2(55)		1,213	19.4(44)	20.9(48)	1,940	7.1 (9)
Lincoln Tr	5,645	8.8 (17)	6.7 (20)		22,919	9.2(50)	3.1(11)	17.4(32)		10,722	19.0(38)	14.9(12)	16,899	8.3 (16)
Little San	1,084	9.1 (24)	9.1 (51)		4,894	4.4 (4)	6.3(53)	13.6 (3)		2,517	15.4 (7)	14.4 (9)	3,004	10.8 (30)
Madison	1,596	10.6 (42)	9.1 (51)		5,532	7.7(32)	5.7(49)	14.2 (7)		2,337	20.0(50)	19.6(42)	3,899	10.3 (27)
Magoffin	499	8.4 (9)	9.4 (55)		2,358	8.4(43)	3.3(14)	25.2(57)		1,280	19.6(47)	22.5(56)	1,498	20.8 (57)
Marshall	667	7.3 (2)	8.8 (48)		2,586	9.6(53)	6.1(52)	15.2(12)		1,185	13.3 (6)	9.5 (2)	1,804	10.1 (26)
Martin	479	10.0 (35)	6.7 (20)		2,158	7.5(28)	3.0 (9)	20.4(51)		1,122	20.0(50)	18.7(37)	1,711	6.7 (6)
Mercer	502	11.0 (46)	5.6 (7)		1,902	9.3(51)	3.4(16)	15.4(17)		860	17.7(23)	17.6(30)	1,361	7.9 (13)
Monroe	309	8.7 (15)	8.7 (46)		1,238	7.7(32)	3.9(24)	15.2(12)		579	16.2(13)	15.9(19)	808	14.6 (46)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)													
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Montgomery	687	10.6 (42)	7.3 (30)	2,743	7.8(35)	2.6 (6)	14.1 (6)	1,312	15.5 (9)	13.0 (4)		2,103	8.1 (14)
Muhlenberg	787	8.4 (9)	7.4 (33)	3,012	4.7 (5)	5.6(48)	14.2 (7)	1,467	15.4 (7)	16.8(25)		1,770	11.6 (33)
N Central	1,443	10.0 (35)	6.7 (20)	5,226	14.3(59)	3.2(12)	27.4(59)	2,200	20.3(53)	23.2(57)		3,637	11.5 (32)
N KY	5,096	10.5 (41)	8.1 (43)	19,711	8.4(43)	4.0(26)	18.3(42)	8,757	18.8(35)	18.4(35)		13,960	12.5 (37)
Oldham	513	12.3 (55)	5.8 (10)	2,077	12.1(57)	2.3 (2)	20.3(50)	959	18.1(29)	17.9(32)		1,281	13.1 (41)
Park DuV	1,208	15.4 (58)	4.7 (1)	4,229	10.6(56)	4.4(33)	15.1(11)	1,934	19.8(49)	15.5(15)		3,109	9.0 (20)
Pennyrile	1,721	10.1 (38)	8.0 (42)	7,114	6.8(20)	4.0(26)	16.6(27)	3,249	17.4(19)	15.5(15)		4,936	14.4 (45)
Pike	1,659	10.1 (38)	7.3 (30)	6,101	3.6 (2)	6.7(55)	17.6(34)	2,754	16.3(14)	20.4(47)		3,878	7.4 (11)
Powell	487	9.9 (31)	4.7 (1)	2,118	6.3(14)	2.3 (2)	18.2(41)	1,102	19.6(47)	20.0(45)		1,687	6.3 (4)
Purchase	3,211	9.6 (30)	7.5 (36)	12,443	7.9(37)	4.6(36)	19.6(47)	5,837	18.6(31)	17.5(28)		8,592	18.1 (53)
Todd	370	11.1 (48)	4.9 (3)	1,455	9.6(53)	4.5(34)	16.4(25)	698	12.9 (4)	11.2 (3)		1,084	18.0 (52)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6F
2002 - 2004 Pediatric Nutrition Surveillance (1)
Kentucky
Comparison of Growth and Anemia Indicators by Local Agency
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Local Agency ^(3,4)														
Local Agency	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)			
	Low		High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)		Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)		Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
WEDCO	1,412	10.8 (44)	5.7 (8)		4,875	7.5(28)	5.0(40)	16.0(23)		2,118	11.9 (1)	13.2 (6)	3,343	12.9 (39)
Whitley	1,720	9.1 (24)	9.1 (51)		7,118	8.2(42)	4.3(32)	19.1(45)		3,421	19.0(38)	21.2(51)	5,285	6.3 (4)
Woodford	297	9.4 (28)	9.1 (51)		1,151	6.4(17)	4.7(37)	15.4(17)		466	12.4 (2)	13.1 (5)	728	13.3 (42)
Unknown Local Agenc	3,339	8.3 (7)	7.0 (27)		3,071	0.7 (1)	9.0(59)	3.8 (1)		92	* *	* *	118	15.3 (47)
Kentucky	99,498	9.7	7.2		367,082	7.6	4.3	17.2		166,051	17.9	17.2	243,357	11.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 1
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
3 Rivers	1,979	19.5 (14)	1,011	3.8 (48)	1,107	2.8 (38)
Allen	765	22.7 (7)	398	4.3 (40)	444	2.7 (42)
Anderson	493	26.6 (4)	250	7.6 (9)	240	4.6 (11)
Barren Riv	9,480	17.4 (22)	4,742	5.1 (25)	4,699	2.9 (35)
Bourbon	651	16.9 (26)	373	5.4 (21)	407	4.2 (16)
Boyd	2,085	14.0 (39)	987	4.8 (28)	1,057	3.0 (34)
Boyle	954	9.6 (53)	465	3.2 (54)	511	1.8 (56)
Bracken	362	19.1 (15)	169	5.9 (17)	177	5.1 (5)
Breathitt	807	7.3 (56)	472	4.4 (36)	511	4.1 (19)
Breckinrid	766	14.8 (36)	402	3.7 (49)	410	2.7 (42)
Buffalo Tr	874	17.3 (23)	456	3.5 (52)	507	2.4 (46)
Bullitt	1,853	18.7 (19)	864	4.5 (34)	957	2.6 (44)
C & Y	1,870	12.9 (47)	918	5.4 (21)	921	2.8 (38)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 2
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Christian	6,254	20.1 (12)	3,676	8.3 (5)	3,081	4.1 (19)
Clark	1,284	17.0 (25)	640	4.4 (36)	622	1.9 (55)
Cumb Val	7,272	12.4 (48)	4,010	4.0 (44)	4,423	2.6 (44)
Estill	807	13.3 (44)	410	4.6 (32)	455	4.0 (21)
Fayette	7,901	35.8 (1)	3,875	12.5 (1)	3,889	6.2 (2)
Fleming	703	12.1 (50)	400	4.5 (34)	441	3.6 (24)
Floyd	2,381	6.7 (57)	1,293	3.9 (45)	1,421	3.1 (32)
Franklin	1,616	24.1 (5)	777	7.3 (14)	688	5.2 (4)
Garrard	686	15.9 (28)	387	3.9 (45)	456	2.9 (35)
Gateway	2,659	13.5 (41)	1,500	4.4 (36)	1,612	3.3 (29)
Green Riv	8,930	15.2 (35)	4,634	4.4 (36)	4,470	2.8 (38)
Greenup	1,326	14.8 (36)	733	5.3 (24)	707	4.2 (16)
Hopkins	1,979	14.0 (39)	966	4.1 (43)	955	2.1 (51)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 3
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Jefferson	18,487	21.9 (8)	8,922	7.7 (7)	8,699	4.7 (10)
Jessamine	1,527	18.8 (18)	834	10.2 (2)	795	6.4 (1)
Johnson	1,327	15.3 (31)	733	4.8 (28)	800	3.4 (27)
Knox	1,858	9.0 (55)	1,027	2.9 (55)	1,107	2.0 (54)
KY River	5,872	10.2 (51)	3,172	4.6 (32)	3,255	3.3 (29)
Lake Cumb	10,016	15.3 (31)	5,412	4.9 (27)	5,740	3.1 (32)
Laurel	2,475	13.4 (42)	1,242	3.5 (52)	1,204	2.4 (46)
Lawrence	813	9.3 (54)	436	5.0 (26)	468	4.3 (15)
Lewis	695	15.3 (31)	362	3.9 (45)	407	2.2 (49)
Lincoln	1,108	15.6 (29)	615	3.7 (49)	555	2.2 (49)
Lincoln Tr	10,040	17.3 (23)	5,454	6.5 (15)	5,335	4.6 (11)
Little San	1,944	17.6 (20)	1,101	4.7 (31)	1,193	2.8 (38)
Madison	2,680	18.9 (16)	1,398	7.6 (9)	1,363	5.1 (5)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Page: 4
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Magoffin	885	6.7 (57)	515	1.6 (58)	562	1.1 (57)
Marshall	1,161	20.3 (11)	626	7.5 (13)	672	5.1 (5)
Martin	831	6.5 (59)	443	2.9 (55)	525	2.1 (51)
Mercer	871	13.4 (42)	438	8.0 (6)	433	4.8 (9)
Monroe	523	17.6 (20)	295	5.4 (21)	274	2.9 (35)
Montgomery	1,245	15.4 (30)	678	8.6 (4)	640	5.3 (3)
Muhlenberg	1,275	10.1 (52)	621	2.4 (57)	717	1.0 (58)
N Central	2,526	16.4 (27)	1,317	5.8 (19)	1,411	3.4 (27)
N KY	9,063	14.1 (38)	5,055	5.9 (17)	4,980	4.0 (21)
Oldham	938	28.9 (3)	511	9.4 (3)	549	4.6 (11)
Park DuV	1,953	18.9 (16)	919	7.7 (7)	913	4.6 (11)
Pennyrile	3,173	22.9 (6)	1,848	7.6 (9)	1,956	5.1 (5)
Pike	2,814	12.4 (48)	1,508	3.6 (51)	1,450	2.3 (48)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7F
2002 - 2004 Pediatric Nutrition Surveillance ⁽¹⁾
Kentucky

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Page: 5
Run Date: 9 / 21 / 05

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Local Agency ^(4,5)						
Local Agency	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Powell	880	13.0 (46)	487	4.3 (40)	544	3.3 (29)
Purchase	5,621	20.1 (12)	3,034	5.8 (19)	3,392	3.8 (23)
Todd	633	20.5 (10)	321	6.2 (16)	363	3.6 (24)
WEDCO	2,363	20.9 (9)	1,224	4.3 (40)	1,264	2.1 (51)
Whitley	3,017	13.3 (44)	1,649	4.8 (28)	1,624	3.5 (26)
Woodford	534	29.0 (2)	291	7.6 (9)	312	4.2 (16)
Unknown Local Agencies	3,350	15.3 (31)	9	* *	19	* *
Kentucky	169,235	17.4	87,305	5.9	88,689	3.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this local agency's rate to other local agencies. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.